


Dispersion Relation and Pulse Shaping of Femtosecond Laser-Driven Transient Pulsed Electromagnetic Fields



Principal Investigators:


P. Bradford, M. Ehret

Expert technical support and physical oversight:

C. Vlachos, M. Krupka, Ph. Korneev, G. Gatti, T. Pisarczyk, J. J. Santos

- 1. Introduction to laser-driven EM discharges**
2. Our work on fast polarisation pulses driven by relativistic lasers
3. Our experiment on ELI-NP

Relativistic laser-solid interactions produce **electromagnetic pulses (EMP)** that propagate in "vacuum" and along the surface of the target



Hot electrons ejected by the laser polarise the target

[A. Brantov *et al.* PRE 2020]

Polarisation pulse propagates over target surface and EM pulse in free space

[K. Quinn *et al.* PRL 2009] [M. Ehret *et al.* PoP submitted]
[S. Kar *et al.* Nat. Comm. 2016]

A return current is drawn when the polarisation wave reaches the ground

[F. Consoli *et al.* HPLSE 2020]

Antenna radiation is emitted when the return current passes along the target support


[J.-L. Dubois *et al.* PRE 2014]
[A. Poyé *et al.* PRE 2015]

The fast polarisation pulse ($t \lesssim 50$ ps)


As hot e^- s cross target-vacuum boundary they excite an EM pulse that propagates in the skin-layer of the target at $v \sim c$

[K. Quinn *et al.* PRL 2009]



$\tau \sim 10$ ps Sommerfeld-type EM surface wave
Low dispersion over cm-distances



[S. Tokita *et al.* Sci. Rep. 2015]




Relativistic e^- s can be entrained within the surface wave



[H. Nakajima *et al.* PRL 2013]

[A. Brantov *et al.* EPJ Web of Conferences 2018]


Electric fields for lensing of charged particle beams



[H. Ahmed *et al.* HPLSE 2017]


The Return Current ($t \gtrsim 100$ ps)

500 fs laser @ $5 \times 10^{18} \text{ Wcm}^{-2} \Rightarrow \sim \text{few kA / tesla in coil}$




[M. Ehret *et al.* PoP submitted]

30 fs laser @ $10^{18} \text{ Wcm}^{-2} \Rightarrow 20 \text{ T coil field}$




[W. Wang *et al.* PoP 2018]

A similar phenomenon is observed at lower intensity ($\lesssim 10^{17} \text{ Wcm}^{-2}$) and longer pulse duration ($t_{\text{las}} \sim \text{ns}$), where the return current can produce quasi-static B-fields




[J. J. Santos *et al.* PoP 2018]

Evidence of multi-kA currents, wire plasma...



[M. Manuel *et al.* APL 2012]



...and long-lived GV/m E-fields




[P. Bradford *et al.* PPCF 2021]

Discharge currents as secondary sources of radiation


Bright THz sources



[N. Bukharskii *et al.* APL 2022]




[Y. Tian *et al.* Nat. Phot. 2017]




[G.-Q. Liao *et al.* PRX 2020]

[N. Bukharskii and Ph. Korneev ArXiV 2022]

GHz EMP





[P. Bradford *et al.* HPLSE 2018]



1. Introduction to laser-driven EM discharges
2. **Our work on fast polarisation pulses driven by relativistic lasers**
3. Our experiment on ELI-NP

Sub- c group velocity and Weibull-like "shape" of fast discharges measured with proton radiography




1. Analytical deflections dominated by radial E-field:

Caustic width => Weibull-like shape

2. **Electrodynamic p+ tracing simulation** => amplitude, $Q = 300 \text{ nC}$


3. Group velocity, $v_g \sim 0.8 - 1 c$



Proton lensing platform provokes interesting questions about the discharge mechanism...

Focusing at coil centre

[M. Ehret *et al.* PoP submitted]



Pulse Characteristics

Rise time, $\tau \sim 10$ ps

FWHM ~ 50 ps

$v_g = 0.8 - 1 c$

$E \sim 10$ GV/m

$B \sim 1 - 10$ T

$I \sim 1 - 10$ kA

Why is the rise and decay $\gg t_{\text{las}}$?

Why is $v_g < c$ on several shots ?



... that are important to answer if we are to develop EM lenses or secondary sources of x-rays, GHz and THz radiation

"Sun-rays" and sub- c group velocities indicative of plasma formation on the wire surface?


Conjecture 1

For a Sommerfeld wave propagating along a solid conductor, have $v_g \sim c$


Sub- c velocity \Rightarrow plasma formation on wire surface?

Conjecture 2

Potential modulations in p+ radiography \Rightarrow surface roughness can modify wave propagation speed and phase?



PIC sims reveal sub- c group velocity of discharge and possible plasma formation on wire




[M. Ehret *et al.* PoP submitted]

Laser drive


$\lambda = 800$ nm,
 $\tau = 1300$ fs,
 $I = 10^{19}$ Wcm $^{-2}$

Surface EM pulse propagates more slowly than the airborne EMP



B-field/current rise-time $\tau \sim 1$ ps is too fast to probe with p+ radiography

Plasma formation on wire and/or directly accelerated e $^{-}$ s?



ChoCoLaTII: Simplified Numerical Modelling of Target Charging

[A. Poyé *et al.* PRE 2018]

Target charging is governed by competition between the hot electron temperature T_e and target potential $\Delta\Phi$


average vel. of electrons that escape the barrier $\Delta\Phi$

$$I_h = e\Omega_b \underbrace{\left(\frac{\eta_\Phi N_h}{V_h}\right)}_{\text{\# density of electrons with energy } > \Delta\Phi} \pi R_h^2 \overbrace{\int_{\Delta\Phi}^{\infty} f_h(\epsilon) v_h d\epsilon}$$

density of electrons with energy $> \Delta\Phi$

Target potential $\Delta\Phi$ can be divided into thermal (from sheath) and electrostatic (from accumulated charge) components


ChoCoLaTII: At high intensity, the rise time of the charge profile can be $\gg t_{\text{las}}$



This is principally due to a long hot e⁻ collisional cooling time

Analytic modelling of EM wave propagating in wire surface plasma suggests we may be able to tune pulse velocity and shape

Plasma Dispersion Relation: Frequency of polarisation pulse envelope related to plasma (n_e , T_e)



We can estimate the **frequency ν** of the pulse by measuring:

plasma (n_e , T_e)


group velocity, v_g

This allows us to discriminate between:

- Expanding surface charge (ChoCoLaTII - long rise-time)
- Sommerfeld wave induced by e^- ejection from target ($\tau_{\text{pulse}} \approx$ laser pulse duration)

Analytic modelling of EM wave propagating in wire surface plasma suggests we may be able to tune pulse velocity and shape

Plasma Dispersion Relation: Frequency of pulse envelope is related to plasma (n_e, T_e)



We can estimate the **frequency ν** of the pulse by measuring:

plasma (n_e, T_e)
group velocity, v_g

This allows us to discriminate between:


- Expanding surface charge (ChoCoLaTII - long rise-time)
- Sommerfeld wave induced by e^- ejection from target ($\tau_{\text{pulse}} \approx \text{laser pulse duration}$)

1. Introduction to laser-driven EM discharges
2. Our work on fast polarisation pulses driven by relativistic lasers
3. **Our experiment on ELI-NP**

ELI-NP is needed to definitively answer the physics of target discharging

We have seen: p+ radiography is limited to $\sim 1-10$ ps resolution

Need **sub-ps resolution** (of order t_{las}) to observe if the current rise time is $\gg t_{\text{las}}$



Quantify impact of E_{las} , t_{las} and focal spot size on the shape (spatial and temporal) of the discharge pulse
(ELI-NP ESSENTIAL)

Focal spot scan $\Rightarrow T_h \stackrel{?}{\Rightarrow} Q$

Study the impact of plasma formation on v_g of polarisation pulse

E_{las} scan (fixed I_{las}) $\stackrel{?}{\Rightarrow}$ impact on Q and T_e via N_h

Study the effect of wire thickness, conductivity and roughness on v_g and phase

t_{las} (fixed I_{las} or fixed E_{las}) $\stackrel{?}{\Rightarrow}$ effect on current and t_{pulse}

Develop pulse trains for selective focusing

Using **optical probing** to diagnose wire currents and plasma conditions at sub-ps resolution

[T. Pisarczyk *et al.* in prep.]

Coil Reference Image




Coil B-field induces polarisation rotation of probe beam

$$E_{\text{probe}} = 10 \text{ mJ}$$


$$\tau_{\text{probe}} \sim 40 \text{ fs}$$

$$\lambda = 800 \text{ nm}$$

Faraday Rotation Scheme



[T. Pisarczyk *et al.* J. Inst, 2019]
[C. Vlachos *et al.* in prep]



LASER



Complex Interferometry

Faraday Rotation

ELI-NP Proposal: First fully-resolved measurement of polarisation pulse and return current

Primary Diagnostics


Interferometry and x-ray spectroscopy to infer plasma n_e and T_e

Faraday rotation measurement of polarisation/return currents

Interferometry:

n_e probe ($2\omega \sim 50\text{mJ}$, 25-200fs)




X-ray spectroscopy: T_e



Test charging model from [Poyé *et al.* PRE 2015]:

Conclusions

There remain important questions about the velocity, amplitude and shape of laser-driven currents...



[N. Bukharskii *et al.* APL 2022]

On ELI-NP:

Our experiment will measure the current evolution at **>10x better temporal resolution** than p+ radiography

We will examine how the discharge current is affected by the **plasma dispersion relation** of the wire

We want to use laser/target parameters to produce **shaped kA current pulses and pulse trains**