

universität freiburg

# High-repetition-rate experiments at ELI-ALPS

Attosecond and Strong-field Physics Group  
Institute of Physics University of Freiburg  
Giuseppe Sansone


**[ELI Ultrafast Science Workshop](#)**

12. February 2026

European XFEL

# Attosecond photoionization delay

Electron passing through a short-range potential accumulates quantum phase




Asymptotic phase difference vs free wave propagation

**Wigner-Smith delay:** 
$$\tau_W = \frac{\partial \eta}{\partial E}$$

E. Wigner, *Phys Rev* **98**, 145 (1955).  
 F. Smith, *Phys Rev* **118**, 349 (1960).  
 L. Argenti *et al.*, *Phys Rev. A* **95** (2017).




J. Vos *et al.*, *Science* **360**, 1326 (2018).  
 M. Huppert *et al.* *PRL* **117**, 093001 (2016).  
 S. Biswas *et al.* *Nature Physics* **16**, 778–783 (2020)  
 H. Ahmadi *et al.*, *Nature Communications* **13**, 1242 (2022).



*Photoemission delays as a sensitive probe*

- Influence of the molecular potential
- Presence of electronic resonances (Shape resonances)
- Influence of electronic correlation effects (shake-up states)
- Coupled nuclear-electronic dynamics <sup>2</sup>

# Attosecond photoionization delays using coincidence spectroscopy






- Measurements under the „same“ experimental conditions
- Angular and energy-resolved information
- Information on the different ionization channels
- Attosecond sources operating at 50 kHz
- Acquisition time (about one week)

J. Ullrich *et al.* J. Phys. B At. Mol. Opt. Phys. 30, 2917–2974 (1997).


J. Ullrich *et al.* Prog. Phys. 66, 1463–1545 (2003).

R. Dörner *et al.* Phys. Rep. 330, 95–192 (2000).


# Attosecond interferometry in CO<sub>2</sub>



## Ion time of flight CO<sub>2</sub> /Ar mixture





## RABBIT spectrogram in CO<sub>2</sub><sup>+</sup>



A. Kamalov et al. PRA **102** (2020).


# Resonant coupling in the cation




$$\Delta E = E(B) - E(C) \approx 1.3 \text{ eV}$$

$$\hbar\omega \approx 1.22 \text{ eV (@1012 nm)}$$


Ion time of flight  $\text{CO}_2/\text{Ar}$  mixture



RABBIT spectrogram in  $\text{CO}_2^+$  B state




# RABBIT: two-pathway photoelectron interferometer




Photoionization leaves the target in an entangled state

$$\Psi(t) = \sum_j \psi_j(t) \Phi_j(t)$$



$$e_{2q+1} = (2q+1) \hbar\omega_{IR} - I_P(B)$$



$$e_{2q-1} = (2q-1) \hbar\omega_{IR} - I_P(B)$$


# 3<sup>rd</sup> - pathway in RABBIT interferometer

- due to the interaction of the IR with the residual ion

➡ 3<sup>rd</sup> pathway arises

$$e_C = (2q+1) \hbar\omega_{IR} - I_P(C) = 2q \hbar\omega_{IR} - I_P(B) \approx e_{2q}$$

$$I_P(B) - I_P(C) = \hbar\omega_{IR}$$



$$e_{2q} = 2q \hbar\omega_{IR} - I_P(B)$$

- IR couples the cationic states without modifying the kinetic energy of the photoelectron

$\Phi_C \Rightarrow \Phi_B$


# 3<sup>rd</sup> path introduces an additional time delay

Total wave function: superposition of photoelectron coupled to **B** cationic state & photoelectron coupled to **C** cationic state.


$$\Psi_n(t) = \Psi_{Bn}(t) \Phi_B(t) + \Psi_{Cn}(t) \Phi_C(t)$$

- Laser induced dipole coupling in the residual ion introduces an additional time delay:

$$\tau_B = \tau_e + \tau_{ion}$$



# Effect of the ionic coupling onto the time delays



- **Close coupling model (CC)** with or without B-C coupling, considers dipole transitions in the residual ion
- **Static exchange (SE)**: single channel picture

➔ Stationary multiphoton R-matrix approach for the equilibrium geometry of the neutral molecule

## Limitations of current setup

- Photon energy range 30-40 eV (mainly limited by driving laser intensity)
- Multi-cycle optical pulses
- Long acquisition time (at 50 kHz)

# HR1 system at ELI-ALPS

HR1: The High Repetition rate laser system provide millijoule level, few-cycle laser pulses at 100 kHz repetition rate, ideally suited for experiments relying on high statistics. The laser systems work in the near infrared, at 1030 nm wavelength.

- ✓ Operation at 100 kHz
- ✓ Few-cycle pulses with CEP tagging
- ✓ Generation of photons up to 100-150 eV
- ✓ Electronic relaxation dynamics and molecular fragmentation
- ✓ Measurement of Auger decay in Krypton (differential photoelectron angular distributions)


Y. Peng *et al* *J. Phys. B: At. Mol. Opt. Phys.* **53** 154004 (2020)

Y. Peng *et al* *Ultrafast Sci.* 2022;2022

T. Csizmadia *et al.* *APL Photonics* **10**, 080803 (2025)

universität freiburg

## Reaction Microscope End Station -REMI-ES






# The team

- **Ioannis Makos**
- David Busto
- Dominik Ertel
- Barbara Merzuk
- Hamed Ahmadi
- Matteo Moioli
- Marvin Schmoll
- Jakub Benda
- Fabio Frassetto
- Luca Poletto
- Claus Dieter Schröter
- Thomas Pfeifer
- Robert Moshhammer
- Zdeněk Mašín
- Serguei Patchkovskii



**Funded by  
the European Union**



**QU-ATTO**

QU-ATTO has received funding from the European Union's Horizon Europe research and innovation programme under the Marie Skłodowska-Curie grant agreement No 101168628

# Thank you for your attention!

Giuseppe Sansone

Attosecond and Strong-Field Physics Group

[giuseppe.sansone@physik.uni-freiburg.de](mailto:giuseppe.sansone@physik.uni-freiburg.de)